

Statement on the need for an arytenoid motion knowledge base

Eric J. Hunter and Laura M. Hunter

National Center for Voice and Speech,
The Denver Center for the Performing Arts, 1245 Champa Street, Denver, CO 80204, USA,

This memo serves as a statement of the need for a compiled knowledge base of arytenoid cartilage movement (i.e., range of motion, speed of motion, etc.). As biomechanical models of the larynx increase in complexity, there is a need to verify their predictions with laboratory data. Fundamental to this are the basics of the arytenoid motion about the cricoarytenoid joint. This memo introduces this need and suggests a course of action. Updates to this memo can be downloaded at <http://www.ncvs.org/ncvs/library/tech>.

Keywords: laryngeal skeleton, cricoarytenoid joint, vocal process, arytenoid motion, range of motion, speed of motion, literature review, cricoid cartilage, arytenoid cartilage, biomechanical models, vocal posturing, vocal fold models, abduction, adduction.

Introduction

Vocal fold abduction/adduction, or posturing, has been hypothesized to be key to vocal onset and self-sustained oscillation (Titze, 1988; Cooke *et al.*, 1997). Variation of adduction in particular has been shown to affect the intensity of the voice (Titze and Sundberg, 1992; Murry *et al.*, 1998), as well as pitch (Hirano and Ohala, 1970; Honda, 1983). Beyond phonation, vocal fold posturing in general is important for ventilation, swallowing, and effort closure of the airway (Kraus *et al.*, 1996).

Numerous studies have attempted to expand this understanding of posturing mechanisms by focusing on the arytenoid cartilage motion around the cricoarytenoid joint (CAJ), as well as the attached muscles causing this motion. Misconceptions about CAJ movement were corrected in a few early studies (*e.g.*, Frable, 1961; von Leden and Moore, 1961; Ardran *et al.*, 1967; Sellars and Sellars, 1983). More recent studies examined details of the CAJ, such as joint axis and stiffness (*e.g.*, Berry *et al.*, 2003; Bryant *et al.*, 1996; Selbie *et al.*, 1998), or arytenoid and vocal fold motion about the CAJ (*e.g.*, Wang, 1998; Hirano *et al.*, 1988).

Hunter & Hunter: Statement on the need for an arytenoid motion knowledge base**The need for a knowledge base**

Notwithstanding the importance of these studies, the results have not yet been combined into a concise, usable knowledge base. Such a knowledge base would be useful, as it would consolidate the data in one location. Perhaps more critical, this consolidation would highlight similarities and differences in methodology (*e.g.*, landmark choice or plane of motion) and resulting measures (*e.g.*, average range of motion). By doing so, best-practice study protocols for such investigations could be identified.

An accurate, thorough understanding of posturing is particularly of interest to researchers who explore CAJ mechanics and their effects on phonation through biomechanical models (*e.g.*, Farley, 1996; Titze, 1996, Hunter, *et al.*, 2004; Kob, 2004). Thus, a knowledge base of arytenoid motion could be used to verify the base assumptions of such models.

Additionally, enhancing the understanding of vocal fold posturing mechanics could advance this and other phonosurgical procedures. It would further the goal of many current laryngeal models (of both phonation and posturing) to lay the foundation for vocal injury prediction (Gunter, 2003; Gunter, 2004). Moreover, as these models improve, they could be used to non-invasively simulate the results of small variations in phonosurgical interventions such as vocal fold medialization (Rosa Mde *et al.*, 2003), a treatment of unilateral vocal fold paralysis in which the vocal fold is artificially postured to set the shape and stiffness so that phonation is improved (Neuman *et al.*, 1994).

Finally, such a knowledge base would draw attention to seemingly unintuitive observations. For example, Hunter *et al.*, 2004, showed a potential physical mechanism behind the observation of Woodson *et al.* (1998) that the cricothyroid muscle did not contribute to vocal fold adduction. Observations like these could then be explored through further laboratory studies or through biomechanical models.

Suggested approach

Because of the complexity of arytenoid cartilage motion, previous studies have used different methodologies to study this motion. One such difference is the choice of landmarks (*e.g.*, mid-membranous vocal fold, CAJ axis point, or vocal process). Therefore, it is proposed by the authors that a compiled knowledge base be built focusing on at least one common landmark to quantify the motion; this common landmark would allow for inter-study comparability. Ideally, the primary landmark would be the vocal process (commonly used to quantify glottal width and glottal length in vocal posturing).

As the knowledge base is compiled, doubtless gaps in research will become evident. Thus, the authors propose specific guidelines for future studies of the CAJ mechanics or movement to facilitate comparisons between studies. First, the authors propose that future studies at least include a direct measurement of, or correlates to, the vocal process (regardless of other landmarks that may be measured). It is true that, although the vocal process is the ideal common landmark,

it is not always easy to pinpoint its exact position (Cooke *et al.*, 1997); thus, it is further proposed that clarification be included on how the vocal process is defined. For example, Hunter and Titze (2004) incorporated the range of potential vocal process definitions by using two nearly identical, commonly used vocal process landmarks: (1) the true anatomical vocal process at the tip of the arytenoid, VP_T ; and (2) the medial vocal fold edge near the vocal process, VP_M .

Further, with the vocal process as the proposed landmark, it is proposed that a compiled knowledge base describe the motion of the vocal process with respect to the cricoid cartilage (after Selbie *et al.*, 1998), as shown in Fig. 1 (x : medio-lateral, rightward positive; y : antero-posterior, anterior or frontward positive; z : inferio-superior, superior or upward positive, after Hunter *et al.*, 2004). Figure 2 illustrates the relation of the arytenoid to the cricoid, with the approximate position of the vocal process labeled as VP.




Figure 1. Coordinate system from vantage point of cricoid cartilage.

Conclusions

As biomechanical models increase in complexity and realism, they can become a more useful tool to study vocal injuries and phonosurgeries. However, before models can be tuned specifically to a subject for testing individual phonosurgeries and/or therapies, the non-geometrical/-anatomical assumptions built into a biomechanical model (*e.g.*, various modeling techniques) must be verified using laboratory data. Therefore, the need for a knowledge base of arytenoid motion ranges (via vocal process motion) was proposed.

Such a knowledge base is essential to the progress and use of biomechanical models of the larynx. Once a usable vocal process range knowledge base is available and posturing models are able to predict the ranges, it will become important to ascertain the speed of posturing motion. Tools such as high-speed stroboscopy using new calibration techniques might be used to obtain these speed measures (Larsson and Hertegard, 2004), allowing researchers to make absolute, rather than relative, measures of abduction/adduction speed. Other potential, new techniques use a prism to capture three-dimensional movement of vocal fold surface dynamics (Dollinger and Berry, 2005); these techniques might be adapted to capture overall vocal fold motion on a lab bench.

It is essential that the researchers design future laryngeal studies with an eye on how their data might be used in the future. First, it is crucial




Figure 2. The right half of the cricoid and the arytenoids. Arrows represent the approximate direction of the adductor/abductor muscles, which attach around the muscular process (M.P.). The vocal process, near the attachment of the vocal ligament, is also labeled (V.P.).

Hunter & Hunter: Statement on the need for an arytenoid motion knowledge base

that studies be designed to allow for inter-study comparisons. Further, as the computational power behind biomechanical models improve, models will become more important for medical and therapeutic purposes; thus, studies should provide data for the use of verifying laryngeal models.

References

- G. M. Ardran and F. H. Kemp. The mechanism of the larynx. I. The movements of the arytenoid and cricoid cartilages. *The British Journal of Radiology* 39 (465):641-654, 1966.
- D. A. Berry, D. W. Montequin, R. W. Chan, I. R. Titze, and H. T. Hoffman. An investigation of cricoarytenoid joint mechanics using simulated muscle forces. *J.Voice* 17 (1):47-62, 2003.
- N. J. Bryant, G. E. Woodson, K. Kaufman, C. Rosen, A. Hengesteg, N. Chen, and D. Yeung. Human posterior cricoarytenoid muscle compartments. Anatomy and mechanics. *Arch.Otolaryngol.Head Neck Surg.* 122 (12):1331-1336, 1996.
- A. Cooke, C. L. Ludlow, N. Hallett, and W. S. Selbie. Characteristics of vocal fold adduction related to voice onsets. *J.Voice* 11 (1):12-22, 1997.
- M. Dollinger and D. A. Berry. Computation of the three-dimensional medial surface dynamics of the vocal folds. *Journal of Biomechanics* 2005.(In Press)
- G. R. Farley. A biomechanical laryngeal model of voice F-0 and glottal width control. *J.Acoust.Soc.Am.* 100 (6):3794-3812, 1996.
- M. A. Frable. Computation of motion at the cricoarytenoid joint. *Arch.Otolaryngology* 73:551-556, 1961.
- H. E. Gunter. A mechanical model of vocal-fold collision with high spatial and temporal resolution. *J.Acoust.Soc.Am.* 113 (2):994-1000, 2003.
- H. E. Gunter. Modeling mechanical stresses as a factor in the etiology of benign vocal fold lesions. *Journal of Biomechanics* 37 (7):1119-1124, 2004.
- M. Hirano, W. Vennard, and J. Ohala. Regulation of register, pitch and intensity of voice. An electromyographic investigation of intrinsic laryngeal muscles. *Folia Phoniatr.(Basel)* 22 (1):1-20, 1970.
- M. Hirano, K. Kiyokawa, and S. Kurita. Laryngeal Muscles and Glottic Shaping. In: *Vocal Physiology: Voice Production, Mechanisms, and Functions*, edited by Osamu Fujimura, NY:Raven Press, Ltd, 1988, p. 49-65.
- K. Honda. Variability analysis of laryngeal muscle activities. In: *Vocal fold physiology: biomechanics, acoustics and phonatory control*, edited by I. R. Titze and Ronald C. Scherer, Denver, CO:The Denver Center for the Performing Arts, 1983, p. 127-137.
- E. J. Hunter, I. R. Titze, and F. Alipour. A three-dimensional model of vocal fold adduction/abduction. *J.Acoust.Soc.Am.* 115 (4):1747-1759, 2004.
- E. J. Hunter and I. R. Titze. NCVS Memo No 03. Dynamics of a three-dimensional model of vocal fold. *NCVS Online Technical Memo* April 2004, 2004. www.ncvs.org/ncvs/library/tech.
- M. Kob and Christoph Butenweg. A finite element model of the interaction between intra- and extralaryngeal muscles. International Conference on Voice Physiology and Biomechanics. 2004.
- D. H. Kraus, M. K. Ali, R. J. Ginsberg, C. J. Hughes, R. F. Orlikoff, V. W. Rusch, M. E. Burt, P. M. McCormack, and M. S. Bains. Vocal cord medialization for unilateral paralysis associated with intrathoracic malignancies. *Journal of Thoracic and Cardiovascular Surgery* 111 (2):334-339, 1996.
- S. Hertegård and H. Larsson. Calibration of high-speed imaging by laser triangulation. *Logopedics Phoniatics Vocology* 29 (4):154-161, 2004.
- T. Murry, Jie Jie Xu, and G. E. Woodson. Glottal configuration associated with fundamental frequency and vocal register. *J.Voice* 12 (1):44-49, 1998.
- T. R. Neuman, A. Hengesteg, R. P. Lepage, K. R. Kaufman, and G. E. Woodson. Three-dimensional motion of the arytenoid adduction procedure in cadaver larynges. *Ann.Otol.Rhinol.Laryngol.* 103 (4 Pt 1):265-270, 1994.

The National Center for Voice and Speech Online Technical Memo, No. 8, 2005

Rev. Feb, 2005

- O. Rosa Mde, J. C. Pereira, M. Grellet, and A. Alwan. A contribution to simulating a three-dimensional larynx model using the finite element method. *J.Acoust.Soc.Am.* 114 (5):2893-2905, 2003.
- W. S. Selbie, L. Zhang, W. S. Levine, and C. L. Ludlow. Using joint geometry to determine the motion of the cricoarytenoid joint. *J.Acoust.Soc.Am.* 103 (2):1115-1127, 1998.
- I. E. Sellars and S. L. Sellars. Cricoarytenoid joint structure and function. *J.Laryngol.Otol.* 97 (11):1027-1034, 1983.
- I. R. Titze. The physics of small-amplitude oscillation of the vocal folds. *J.Acoust.Soc.Am.* 83 (4):1536-1552, 1988.
- I. R. Titze and J. Sundberg. Vocal intensity in speakers and singers. *J.Acoust.Soc.Am.* 91 (5):2936-2946, 1992.
- I. R. Titze. Coupling of neural and mechanical oscillators in control of pitch, vibrato, and tremor. In: *Vocal fold physiology: controlling complexity and chaos*, edited by P. J. Davis and Neville H. Fletcher, San Diego, CA:Singular Publishing Group, 1996, p. 47-62.
- E. von Leden and P. Moore. The mechanics of the cricoarytenoid joint. *Arch.Otolaryngol.* 73:541-550, 1961.
- R. C. Wang. Three-dimensional analysis of cricoarytenoid joint motion. *Laryngoscope* 108 (4 Pt 2 Suppl 86):1-17, 1998.
- G. E. Woodson, M. P. Murry, V. Schweizer, A. Hengesteg, N. Chen, and D. Yeung. Unilateral cricothyroid contraction and glottic configuration. *J.Voice* 12 (3):335-339, 1998.

Acknowledgements

This work is part of a research program entitled, "A Computational Tool for Simulation of Phonosurgical Procedures", supported by NIDCD grant DC006801 from the National Institutes of Health. Thanks to the Denver Center for the Performing Arts and their continued support as the NCVS Denver site host.

Copyright

The images and text are open to use by the public as a service of the National Center for Voice and Speech. The images and text enclosed in this memo and accompanying this memo are open to use by the public as a service of the National Center for Voice and Speech. However, we ask the reader to respect the time and effort put into this manuscript and research. If the text or images are used, the user agrees to reference to this document and the National Center for Voice and Speech

Revisions

1.0 Eric Hunter, main content and formatted (Feb-2005)